

Definitions and Formulas for Advanced Mathematics

LOGIC		ALGEBRA	
$a \rightarrow b$	a implies b	$i = \sqrt{-1}$	imaginary unit
$a \leftrightarrow b$	a if and only if b	\bar{z}	complex conjugate of z
$a \wedge b$	a and b	A^{-1}	inverse of matrix A
$a \vee b$	a or b	\vec{v}	vector v
$\sim a$	not a		
$A \cup B$	A union B		
$A \cap B$	A intersect B		
\bar{A}	complement of A		
U	universal set		
{ }	empty set		

GEOMETRY


$$S = 4\pi r^2$$
 surface area of a sphere

$$V = \frac{4}{3}\pi r^3$$
 volume of a sphere


\sim is similar to

\cong is congruent to

Congruent Angles




Congruent Sides



Parallel Lines






Hero's or Heron's Formula



$$\text{Area} = \sqrt{s(s - a)(s - b)(s - c)}$$

$$\text{where } s = \frac{a + b + c}{2}$$

GEOMETRY (continued)



Eccentricity of a Conic

$$e = \frac{c}{a}$$

Directrices of a Conic

$$x = \pm \frac{a}{e} = \pm \frac{a^2}{c}$$

TRIGONOMETRY

$$\sin(\theta_1 \pm \theta_2) = \sin \theta_1 \cos \theta_2 \pm \cos \theta_1 \sin \theta_2$$


$$\cos(\theta_1 \pm \theta_2) = \cos \theta_1 \cos \theta_2 \mp \sin \theta_1 \sin \theta_2$$

$$\tan(\theta_1 \pm \theta_2) = \frac{\tan \theta_1 \pm \tan \theta_2}{1 \mp \tan \theta_1 \tan \theta_2}$$

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$



Law of Sines

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

Law of Cosines

$$c^2 = a^2 + b^2 - 2ab \cos C$$

STATISTICS

$$\text{standard deviation of a sample mean} = \frac{\sigma}{\sqrt{N}}$$

NOTES FOR ADVANCED MATHEMATICS TEST

In this examination, assume all functions are real valued functions unless otherwise noted.

In this examination, diagrams may not be drawn to scale.

In this examination, assume all geometry problems imply the use of Euclidean geometry unless otherwise noted.